

For patients who take Tablet 'F' (levonorgestrel) as an emergency contraceptive

■ Indications

Tablet 'F' (levonorgestrel 1.5 mg) is an emergency contraceptive that helps to prevent pregnancy after birth control failure or unprotected sexual intercourse.

■ Dosage and administration

Take one tablet as soon as possible **within 72 hours after unprotected** sexual intercourse.

This tablet can be taken at any time of the day but should be taken **as soon as possible after unprotected** sexual intercourse.



Levonorgestrel is a white-colored tablet and is available in blister packs containing one tablet (single dose).

Do not take levonorgestrel if:

- * You have ever developed hypersensitivity to any of the ingredients of this medicine.
- * You have a severe liver disorder.
- * You are pregnant.

Talk to your doctor if:

- * You have liver disorder.
- * You have heart disease.
- * You have kidney disease.

Precautions

- This medicine cannot always prevent pregnancy.
- This medicine is an emergency contraceptive that helps to prevent pregnancy after birth control failure or unprotected sexual intercourse and should not be used as a regular method of contraception like ordinary contraceptives.

Directions after use

- If you vomit within two hours after taking the medicine, contact your doctor immediately.
- As you have the possibility to be pregnant even after taking this medicine, you should take appropriate contraceptive measures.
- As it may be difficult to distinguish between abnormal genital bleeding or bleeding in early pregnancy and normal menstruation after taking this medicine, you should visit your doctor and get medical care.
- The active ingredient of this medicine is excreted via your breast milk. Therefore, if you are breastfeeding, avoid breastfeeding for 24 hours after taking this medicine.
- After taking this medicine, you might have some bleeding like menstruation (withdrawal bleeding). You might also have symptoms that appear during menstruation, such as abnormal uterine bleeding, headache, nausea, fatigue, and drowsiness. If these symptoms persist, please consult your doctor.
- After taking this medicine, there may be temporary changes in your menstrual cycle or bleeding (the amount and number of days of bleeding).

You may take an oral contraceptive for regular birth control after emergency contraception.